

SYSTOLIC VOLUME AND MINIMAL ENTROPY OF ASPHERICAL MANIFOLDS

STÉPHANE SABOURAU

ABSTRACT. We establish isosystolic inequalities for a class of manifolds which includes the aspherical manifolds. In particular, we relate the systolic volume of aspherical manifolds first to their minimal entropy, then to the algebraic entropy of their fundamental groups.

CONTENTS

1. Introduction	1
2. Pseudomanifolds, relative systole and relative entropy	6
3. Minimal entropy of geometric cycles representing aspherical manifolds	9
4. Entropy of extremal geometric cycles	15
5. Relative systolic volume and relative minimal entropy	17
6. Relative systolic volume and algebraic entropy	19
References	22

1. INTRODUCTION

Let M be a nonsimply connected closed n -manifold endowed with a Riemannian metric g . The (homotopy) systole of (M, g) , denoted $\text{sys}(M, g)$, is defined as the length of the shortest noncontractible loop in M . The optimal systolic volume of M , denoted $\sigma(M)$, is defined as

$$\sigma(M) = \inf_g \frac{\text{vol}(M, g)}{\text{sys}(M, g)^n} \quad (1.1)$$

where g runs over the space of all metrics on M .

1991 *Mathematics Subject Classification.* 53C20, 53C23.

Key words and phrases. systole, systolic volume, isosystolic inequality, volume entropy, minimal entropy, algebraic entropy.

Current address: Department of Mathematics, University of Pennsylvania, 209 South 33rd Street, Philadelphia, PA 19104-6395, USA.

E-mail: `sabourau@lmpm.univ-tours.fr`.

The study of the systolic volume constitutes the heart of systolic geometry. We refer the reader to the expository texts [12], [19], [20] and [14] and the references therein for an account on the subject.

In 1983, M. Gromov proved the following fundamental systolic inequality whose converse for orientable manifolds was established by I. Babenko in [1].

Theorem 1.1 ([18]). *There exists a positive constant C_n such that every n -dimensional essential manifold M satisfies*

$$\sigma(M) \geq C_n. \quad (1.2)$$

Recall that a closed manifold M is said to be essential if there is a map f from M into a $K(\pi, 1)$ such that $f_*([M]\overline{\gamma}) \neq 0$ in $H_n(\pi; \overline{\gamma})$ where $\overline{\gamma} = \mathbb{Z}$ if M is orientable or \mathbb{Z}_2 otherwise. In particular, aspherical manifolds are essential.

This result extends earlier works on surfaces (see the references mentioned above). Note that the precise value of the optimal systolic volume of essential manifolds is known only for the 2-torus (C. Loewner, unpublished, [20]), the projective plane (P.M. Pu, [29]) and the Klein bottle (C. Bavard, [10], [30]).

The systolic inequality (1.2) can be improved by taking into account the topology of the manifold. For instance, closed surfaces of large genus have a large systolic volume (see [18, 5.3]). More generally, M. Gromov showed that the systolic volume of a closed manifold M is bounded from below in terms of its simplicial volume $||M||$, its simplicial height $h(M)$ and, in the aspherical case, the sum of its Betti numbers $b(M) = \sum_i b_i(M)$ (see [19, 3.C] for precise definitions). Specifically, we have

Theorem 1.2 ([18, 6.4.C'' & D'], [19, 3.C] and also [20, 4.46]). *Every n -dimensional closed manifold M satisfies*

$$\sigma(M) \geq C_n \frac{||M||}{\log^n(1 + ||M||)}, \quad (1.3)$$

$$\sigma(M) \geq C_n \frac{h(M)}{\exp(C'_n \sqrt{\log h(M)})}. \quad (1.4)$$

Furthermore, if M is aspherical, we have

$$\sigma(M) \geq C_n \frac{b(M)}{\exp(C'_n \sqrt{\log b(M)})}. \quad (1.5)$$

Here, the positive constants C_n and C'_n depend only on the dimension of M .

Note that inequality (1.5) is a consequence of (1.4) since $h(M) \leq b(M)$ for aspherical manifolds.

In [1], [3] and [4], I. Babenko compared the optimal systolic volumes of the manifolds with the same fundamental group. Namely, he proved that the optimal systolic volume is a homotopy invariant (see [1]). He also showed that in a large number of cases, the optimal systolic volume of a manifold depends only on its homology class in the corresponding Eilenberg-MacLane space (see [3] and [4] for a more precise statement).

In this article, we prove two other systolic inequalities for aspherical manifolds using, on the one hand, the minimal entropy and, on the other hand, the algebraic entropy of the fundamental groups. Loosely speaking, we show that aspherical manifolds with “complicated” fundamental groups have large systolic volume.

In order to state our first result, we need to introduce some definitions.

Given a closed n -dimensional manifold (M, g) , denote by (\tilde{M}, \tilde{g}) the universal Riemannian covering. Fix $x_0 \in M$ and a lift $\tilde{x}_0 \in \tilde{M}$ of x_0 .

The volume entropy (or asymptotic volume) of (M, g) is defined as

$$\text{Ent}(M, g) = \lim_{R \rightarrow +\infty} \frac{\log(\text{vol}_{\tilde{g}} B(\tilde{x}_0, R))}{R} \quad (1.6)$$

where $\text{vol}_{\tilde{g}} B(\tilde{x}_0, R)$ is the volume of the ball centered at \tilde{x}_0 with radius R in \tilde{M} . Since M is compact, the limit in (1.6) exists and does not depend on the point $\tilde{x}_0 \in \tilde{M}$ (see [28]). This asymptotic invariant describes the exponential growth rate of the volume on the universal covering.

Define the minimal entropy of M as

$$\text{MinEnt}(M) = \inf_g \text{Ent}(M, g) \text{vol}(M, g)^{\frac{1}{n}} \quad (1.7)$$

where g runs over the space of all metrics on M .

M. Gromov showed in [17] that the simplicial volume is related to the minimal entropy. More precisely, there exists a positive constant C_n such that every closed n -manifold M with simplicial volume $||M||$ satisfies

$$\text{MinEnt}(M)^n \geq C_n ||M||. \quad (1.8)$$

When M admits a negatively curved locally symmetric metric g_0 , it has been proved by A. Katok in [22] (for $n = 2$) and by G. Besson, G. Courtois and S. Gallot in [11] (for $n \geq 3$) that

$$\text{MinEnt}(M)^n = \text{Ent}(g_0)^n \text{vol}(g_0). \quad (1.9)$$

I. Babenko showed in [1] that both the optimal systolic volume and the minimal volume entropy are homotopy invariants of manifolds.

The first result of this article shows how these two invariants are related. More precisely, we prove the following partial generalization of inequality (1.3) in Theorem 1.2.

Theorem 1.3. *Let $\Phi : M \longrightarrow K$ be a degree 1 map between two oriented closed manifolds of the same dimension $n \neq 3$ such that K is a $K(\pi_1(M), 1)$ -space. Then, there exists a positive constant C_n depending only on n such that*

$$\sigma(M) \geq C_n \frac{\text{MinEnt}(M)^n}{\log^n(1 + \text{MinEnt}(M))}. \quad (1.10)$$

If M is aspherical, we can take the identity map for Φ . This immediately leads to the following result.

Corollary 1.4. *Every orientable aspherical closed manifold of dimension n different from 3 satisfies the systolic inequality (1.10).*

Theorem 1.3 is more general than Corollary 1.4. Indeed, the connected sum of an aspherical manifold satisfying the hypothesis of Theorem 1.3 with a simply connected orientable closed manifold nonhomeomorphic to the sphere again satisfies the hypothesis of Theorem 1.3 but is not aspherical.

Inequality (1.10) combined with (1.8) implies inequality (1.3). Thus, Theorem 1.3 sharpens Theorem 1.2 (1.3) for aspherical manifolds of dimension different from 3, but it does not apply to arbitrary closed manifolds as Theorem 1.2 (1.3) does. It is an open question whether inequality (1.10) holds for every closed manifold.

The restriction on the dimension of M in Theorem 1.3 is due to technical reasons: in the proof, we use a construction of I. Babenko [4] which relies on obstruction theory and requires that n is different from 3 (see Section 3 for further explanation).

Let us now introduce the algebraic entropy of a finitely generated group.

Let Σ be a finite generating set of a group Γ . The algebraic (or word) length of an element $\alpha \in \Gamma$ with respect to Σ is noted $|\alpha|_\Sigma$. It is defined as the smallest integer $k \geq 0$ such that $\alpha = \alpha_1 \dots \alpha_k$ where $\alpha_i \in \Sigma \cup \Sigma^{-1}$. By definition, the neutral element e is the only element of Γ with null algebraic length.

The algebraic entropy of Γ with respect to Σ is defined as

$$\text{Ent}_{alg}(\Gamma, \Sigma) = \limsup_{R \rightarrow +\infty} \frac{\log(N_\Sigma(R))}{R}. \quad (1.11)$$

where $N_\Sigma(R) = \text{card}\{\alpha \in \Gamma \mid |\alpha|_\Sigma \leq R\}$ is the cardinal of the R -ball of $(\Gamma, |\cdot|_\Sigma)$ centered at its origin. If Γ is the fundamental group of a closed polyhedron, then the sup-limit is a limit (see [28]).

Define the minimal algebraic entropy of Γ as

$$\text{Ent}_{alg}(\Gamma) = \inf_{\Sigma} \text{Ent}_{alg}(\Gamma, \Sigma) \quad (1.12)$$

where Σ runs over the space of all generating sets of Γ .

The second result of this article shows how the systolic volume of some manifolds is related to the algebraic entropy of their fundamental groups. More precisely, we have

Theorem 1.5. *Let $\Phi : M \longrightarrow K$ be a degree 1 map between two n -dimensional oriented closed manifolds such that K is a $K(\pi_1(M), 1)$ -space. Then, there exists a positive constant C_n depending only on n such that*

$$\sigma(M) \geq C_n \frac{\text{Ent}_{alg}(\pi_1(M))}{\log(1 + \text{Ent}_{alg}(\pi_1(M)))}. \quad (1.13)$$

Note that the dimension n can be equal to 3 in this result. As previously, we have the following particular result.

Corollary 1.6. *Every n -dimensional orientable aspherical closed manifold satisfies the systolic inequality (1.13).*

As shown in Example 6.5, inequality (1.13) does not hold for every orientable closed manifold: extra topological conditions are required, as in Theorem 1.5.

The constants C_n we obtain in inequalities (1.10) and (1.13) can be explicitly computed. Theorems 1.3 and 1.5 are consequences of more general results - namely, Theorems 5.1 and 6.1 - stated in terms of relative systole and relative entropy (see Section 2 for a definition of these notions). Note that their proofs are not based on filling invariant estimates.

Let us now mention some other related works, though of a quite different nature, on universal systolic inequalities.

Upper bounds on the optimal systolic volume of the connected sums and the cyclic coverings of manifolds can be found in [13] for surfaces and in [5] in higher dimensions.

The (homotopy) systole can be replaced by stable or conformal systoles of any dimension. In this case, multiplicative relations in the cohomology ring lead to systolic inequalities (see [18], [20], [8], [9], [24], [23], [7], [25] for more precise statements and related results). Note that some of these stable/conformal systolic inequalities are sharp.

In contrast with these results, unstable higher dimensional systoles typically satisfy no nontrivial systolic inequalities (see [16], [27] and [2] for the most general and recent results).

This article is organized as follows. The definition of pseudomanifolds and the notions of relative systole and relative entropy are presented in Section 2. In Section 3, we show that under some topological conditions, the relative minimal entropy of geometric cycles representing a manifold is not less than the relative minimal entropy of this manifold. The proof of this comparison result makes use of surgery and extension procedures and relies on I. Babenko's constructions. In Section 4, we introduce regular geometric cycles whose existence and main properties have been established by M. Gromov. Then, we bound from above their relative minimal entropy in terms of their relative systolic volume. Combining the results of Sections 3 and 4, we derive in Section 5 a lower bound on the relative systolic volume of some manifolds in terms of their relative minimal entropy. In Section 6, we show that under some topological conditions, the relative systolic volume of manifolds can be bounded from below in terms of the algebraic entropy of their fundamental group.

2. PSEUDOMANIFOLDS, RELATIVE SYSTOLE AND RELATIVE ENTROPY

In this section, we recall the definition of pseudomanifolds and introduce the notions of relative systole and relative entropy.

The definition of manifolds appears too restrictive in the study of the systolic volume. Instead, we will consider Riemannian polyhedra, i.e., polyhedra endowed with a piecewise linear Riemannian metric (see [1, §2] for a more precise definition). Unless stated otherwise, all polyhedra will be finite. By definition, a finite n -dimensional polyhedron X is said to be orientable if $H_n(X; \mathbb{Z}) \simeq \mathbb{Z}$.

The class of closed orientable n -dimensional pseudomanifolds plays an important role in our study, especially in relation to inequalities (4.3) and (4.4).

Definition 2.1 ([31]). A closed n -dimensional pseudomanifold is a finite simplicial complex X such that

- (1) every simplex of X is a face of some n -simplex of X ;
- (2) every $(n-1)$ -simplex of X is the face of exactly two n -simplices of X ;
- (3) given two n -simplices s and s' of X , there exists a finite sequence $s = s_1, s_2, \dots, s_m = s'$ of n -simplices of X such that s_i and s_{i+1} have an $(n-1)$ -face in common.

The n^{th} homology group $H_n(X; \mathbb{Z})$ of a closed n -dimensional pseudomanifold is either isomorphic to \mathbb{Z} or trivial (see [31]).

Lemma 2.2. *Every closed n -dimensional pseudomanifold admits a cellular decomposition with exactly one n -cell.*

Proof. Consider a closed n -pseudomanifold X with its simplicial structure. There exists a tree T in X whose vertices agree with the barycenters of the n -simplices of X and edges are segments joining the barycenters of adjacent n -simplices. Here, we say that two n -simplices are adjacent if they have an $(n-1)$ -face in common. The $(n-1)$ -simplices of X which do not intersect T form an $(n-1)$ -dimensional complex X' . It is possible to define a cellular complex homeomorphic to X by gluing an n -cell to X' . \square

Let us now extend the classical definitions of the systole and the entropy.

Let X be a finite n -dimensional polyhedron endowed with a piecewise Riemannian metric g . Let $\psi : \pi_1(X) \rightarrow \pi$ be a group homomorphism with kernel $H \triangleleft \pi_1(X)$. Denote by ψ_{\sharp} the map induced by ψ between the conjugacy classes.

The ψ -systole of (X, g) , denoted $\text{sys}_{\psi}(X, g)$, is defined as the length of the shortest loop of X whose image by ψ_{\sharp} is nontrivial.

The ψ -systolic volume of (X, g) is defined as

$$\sigma_{\psi}(X, g) = \frac{\text{vol}(X, g)}{\text{sys}_{\psi}(X, g)^n} \quad (2.1)$$

and the optimal ψ -systolic volume of X is defined as

$$\sigma_{\psi}(X) = \inf_g \sigma_{\psi}(X, g) \quad (2.2)$$

where g runs over the space of all piecewise linear Riemannian metrics on X . Here, the volume of an n -dimensional Riemannian polyhedron (X, g) is defined as the sum of the n -volumes of all the n -simplices of X . Thus, the n -volume agrees with the n -dimensional Hausdorff measure. Note that the optimal ψ -systolic volume of X depends rather on the kernel H of ψ than on the homomorphism ψ itself. We clearly have

$$\sigma_{\psi}(X) \leq \sigma(X). \quad (2.3)$$

Let $p : X_H \rightarrow X$ be the covering corresponding to the normal subgroup $H \triangleleft \pi_1(X)$. By definition, the deck transformation group of X_H , noted $\text{Aut}(X_H)$, agrees with $\pi_1(X)/H$ and $\pi_1(X_H)$ is isomorphic to H . The lift of g to X_H is noted \bar{g} . Fix x_0 in X and a lift \bar{x}_0 in X_H .

The volume entropy of X relative to H (or H -entropy) is defined as

$$\text{Ent}_H(X, g) = \lim_{R \rightarrow +\infty} \frac{1}{R} \log(\text{vol}_{\bar{g}} B(\bar{x}_0, R)) \quad (2.4)$$

where $\text{vol}_{\bar{g}} B(\bar{x}_0, R)$ is the volume of the \bar{g} -ball centered at \bar{x}_0 of radius R in X_H . Since X is compact, the limit in (2.4) exists and does not depend on the point $\bar{x}_0 \in X_H$ (see [28]). When H is trivial, the H -entropy agrees with the classical volume entropy. Furthermore, if $H_1 \triangleleft H_2 \triangleleft \pi_1(M)$, then $\text{Ent}_{H_1}(X, g) \geq \text{Ent}_{H_2}(X, g)$. In particular, one has $\text{Ent}_H(X, g) \leq \text{Ent}(X, g)$.

The minimal H -entropy of X is defined as

$$\text{MinEnt}_H(X) = \inf_g \text{Ent}_H(X, g) \text{vol}(X, g)^{\frac{1}{n}} \quad (2.5)$$

where g runs over the space of all piecewise linear Riemannian metrics on X . When H is trivial, the minimal H -entropy agrees with the classical minimal entropy.

The H -entropy can also be defined by using the following result.

Lemma 2.3. *Let X be a closed polyhedron endowed with a piecewise linear Riemannian metric g and $\psi : \pi_1(X) \longrightarrow \pi$ be a group homomorphism with kernel H . Then,*

$$\text{Ent}_H(X, g) = \lim_{T \rightarrow +\infty} \frac{\log(P_H(T))}{T} \quad (2.6)$$

where $P_H(T)$ is the number of classes in $\text{Aut}(X_H) \simeq \pi_1(X, x_0)/H$ which can be represented by loops of length at most T based at some fixed point x_0 .

Proof. The group $\Gamma := \text{Aut}(X_H) \simeq \pi_1(X, x_0)/H$ acts on X_H by isometries. The orbit of \bar{x}_0 by Γ is noted $\Gamma.\bar{x}_0$ where \bar{x}_0 is a lift of x_0 in X_H . Consider a fundamental domain Δ for the action of Γ containing \bar{x}_0 . Denote by D the diameter of Δ . The cardinal of $\Gamma.\bar{x}_0 \cap B(\bar{x}_0, R)$ is bounded from above by the number of translated fundamental domains $\gamma.\Delta$ contained in $B(\bar{x}_0, R + D)$ and bounded from below by the number of translated fundamental domains $\gamma.\Delta$ contained in $B(\bar{x}_0, R)$. Therefore, we have

$$\frac{\text{vol}(B(\bar{x}_0, R))}{\text{vol}(X, g)} \leq \text{card}(\Gamma.\bar{x}_0 \cap B(\bar{x}_0, R)) \leq \frac{\text{vol}(B(\bar{x}_0, R + D))}{\text{vol}(X, g)}. \quad (2.7)$$

Take the log of these terms, multiply them by $\frac{1}{R}$ and let R go to infinity. The left-hand term and the right-hand term both yield $\text{Ent}_H(X, g)$. The result follows since $P'_H(R) = \text{card}(\Gamma.\bar{x}_0 \cap B(\bar{x}_0, R))$. \square

3. MINIMAL ENTROPY OF GEOMETRIC CYCLES REPRESENTING ASPHERICAL MANIFOLDS

In this section, we compare the relative minimal entropy of some manifolds with the relative minimal entropy of their representing geometric cycles.

Let π be a discrete group and $K(\pi, 1)$ be an Eilenberg-MacLane space. Let h be a homology class in $H_n(\pi; \mathbb{Z}) := H_n(K(\pi, 1); \mathbb{Z})$. A geometric cycle representing the homology class h is a map $\Psi : X \rightarrow K(\pi, 1)$ from a closed oriented n -dimensional Riemannian pseudomanifold (X, g) such that $\Psi_*[X] = h$ where $[X]$ is the fundamental class of X . The map Ψ induces a homomorphism $\psi : \pi_1(X) \rightarrow \pi$ between the fundamental groups, whose kernel is noted H .

The minimal entropy of $h \in H_n(\pi; \mathbb{Z})$ is defined as

$$\text{MinEnt}(h) = \inf_X \text{MinEnt}_H(X) \quad (3.1)$$

where $X = (X, \Psi, g)$ runs over all the geometric cycles representing h .

Theorem 3.1. *Let π be a discrete group. Let $\Phi : M \rightarrow K$ be a degree 1 map between two oriented closed manifolds of the same dimension $n \neq 3$ such that K is a $K(\pi, 1)$ -space. Then,*

$$\text{MinEnt}_G(M) = \text{MinEnt}(\Phi_*[M]) \quad (3.2)$$

where G is the kernel of the homomorphism $\phi : \pi_1(M) \rightarrow \pi$ induced by Φ and $[K]$ is the fundamental class of K .

Remark 3.2. Under the assumptions of Theorem 3.1, we have $\Phi_*[M] = [K]$. We can apply again Theorem 3.1 to the identity map of K . This yields

$$\text{MinEnt}_G(M) = \text{MinEnt}(K). \quad (3.3)$$

Remark 3.3. Every homomorphism $\phi : \pi_1(M) \rightarrow \pi$ induces a map $\Phi : M \rightarrow K$ (uniquely defined up to homotopy) whose homomorphism induced between the fundamental groups agrees with ϕ . Thus, if M is aspherical, we can take $\pi = \pi_1(M)$, $K = K(\pi_1(M), 1) = M$ and Φ the map induced by the identity homomorphism on $\pi_1(M)$. In this case, we obtain

$$\text{MinEnt}(M) = \text{MinEnt}([M]). \quad (3.4)$$

Remark 3.4. Under the hypotheses of Theorem 3.1, the group π is necessarily a quotient of $\pi_1(M)$ (see Lemma 3.7).

Let X be a closed n -dimensional polyhedron and $\psi : \pi_1(X) \rightarrow \pi$ be a group homomorphism with kernel H . An H -extension of X is a closed n -dimensional polyhedron X' obtained by attaching to X a

finite number of k -cells with $2 \leq k \leq n-1$ along simplicial maps $h : S^{k-1} \longrightarrow X'^{(k-1)}$ into the $(k-1)$ -skeleton of X' , with $[h] \in H$ when $k=2$. In particular, an H -extension X' of X contains X and may be written as

$$X' = X \bigcup_{k=2}^{n-1} \bigcup_{i_k=1}^{p_k} B_{i_k}^k. \quad (3.5)$$

The group $\pi_1(X')$ is isomorphic to a quotient of $\pi_1(X)$ by a normal subgroup of H . Thus, the map ψ induces a homomorphism $\psi' : \pi_1(X') \longrightarrow \pi$ whose kernel H' is a quotient of H .

The inclusion map $i : X \hookrightarrow X'$ induces an isomorphism between $H_n(X; \mathbb{Z})$ and $H_n(X'; \mathbb{Z})$. In particular, if X is an orientable polyhedron, then the same holds for X' . In this case, the degree of the inclusion map $i : X \hookrightarrow X'$ equals 1 (i.e., $i_*[X] = [X']$ in $H_n(X'; \mathbb{Z})$ for corresponding orientations). Recall that the degree of a map f between two oriented n -dimensional polyhedra X and X' is defined as the unique integer p such that $f_*[X] = p[X']$ in $H_n(X'; \mathbb{Z})$. When H is trivial, the space X' is simply called an extension of X .

A simplicial map $f : X \longrightarrow Y$ between two finite simplicial polyhedra is said to be n -monotone if the preimage of every open n -simplex of Y is either an open n -simplex of X or an empty set.

The following result is due to I. Babenko (see [1, §2]).

Lemma 3.5. *Let X_i , $i = 1, 2$, be two closed n -dimensional polyhedra and $\psi_i : \pi_1(X_i) \longrightarrow \pi$ be two group homomorphisms with kernel H_i . Assume that there exists an n -monotone map $f : X_1 \longrightarrow X_2$ such that $\psi_1 = \psi_2 \circ f_*$. Then,*

$$\text{MinEnt}_{H_1}(X_1) \leq \text{MinEnt}_{H_2}(X_2) \quad (3.6)$$

Proof. Fix $\varepsilon > 0$ and a metric g on X_2 . From [1, §2] (see also [4, Lemme 3.1]), there exists a metric g_ε on X_1 such that $f : (X_1, g_\varepsilon) \longrightarrow (X_2, g)$ is nonexpanding and $\text{vol}(X_1, g_\varepsilon) \leq \text{vol}(X_2, g) + \varepsilon$. Since $\psi_1 = \psi_2 \circ f_*$, we have $f_*^{-1}(H_2) = H_1$. Thus, f_* induces an injective homomorphism between $\pi_1(X_1)/H_1$ and $\pi_1(X_2)/H_2$. Moreover, if γ is a loop of X_1 based at some fixed point x_0 , its image $f(\gamma)$ is a loop of X_2 based at $f(x_0)$ such that $\text{length}(f(\gamma)) \leq \text{length}(\gamma)$. Therefore, $P_{H_1}(T) \leq P_{H_2}(T)$. This implies that $\text{Ent}_{H_1}(X_1, g_\varepsilon) \leq \text{Ent}_{H_2}(X_2, g)$ from Lemma 2.3. The result follows. \square

The following result is an immediate consequence of Lemma 3.5.

Lemma 3.6. *Every H -extension X' of X satisfies*

$$\text{MinEnt}_{H'}(X') = \text{MinEnt}_H(X) \quad (3.7)$$

Proof. The inclusion map $i : X \hookrightarrow X'$ is n -monotone and $\psi = \psi' \circ i_*$. Therefore, from Lemma 3.5, we have $\text{MinEnt}_H(X) \leq \text{MinEnt}_{H'}(X')$.

Given a metric g on X , we extend g to the H -extension X' of X by induction on k so that each cell $B_{i_k}^k$ has the geometry of a long cylinder over its basis capped on its top (see [4, Lemma 3.5] for a more precise construction). Denote by g' the metric on X' so-obtained. If the cylinders defining the geometry of the $B_{i_k}^k$'s are long enough, every loop γ' of X' based at some fixed point x_0 of X can be homotoped in X' into a loop γ of X with the same based point such that $\text{length}(\gamma) \leq \text{length}(\gamma')$. Since the inclusion map induces an isomorphism between $\pi_1(X)/H$ and $\pi_1(X')/H'$, we have $P_{H'}(T) \leq P_H(T)$. We deduce from Lemma 2.3 that $\text{Ent}_{H'}(X', g') \leq \text{Ent}_H(X, g)$. Hence, $\text{MinEnt}_{H'}(X') \leq \text{MinEnt}_H(X)$ since the n -volume of the cells attached to X is null. \square

We can now prove Theorem 3.1 when $n \geq 4$. The case $n = 2$ is treated at the end of this section.

Let $\Phi : M \rightarrow K$ be a degree 1 map between two oriented closed manifolds of the same dimension $n \geq 4$ such that K is a $K(\pi, 1)$ -space. Denote by $\phi : \pi_1(M) \rightarrow \pi$ the homomorphism induced by Φ between the fundamental groups. Let $\Psi : X \rightarrow K$ be a geometric cycle representing $\Phi_*[M] = [K]$ in homology. By definition, the map Ψ is of degree 1. Denote by $\psi : \pi_1(X) \rightarrow \pi$ the homomorphism induced by Ψ . The fundamental groups of M and X are related to π as follows.

Lemma 3.7. *The homomorphisms $\phi : \pi_1(M) \rightarrow \pi$ and $\psi : \pi_1(X) \rightarrow \pi$ are epimorphisms.*

Proof. Let $p : \overline{K} \rightarrow K$ be the covering corresponding to the subgroup $\psi(\pi_1(X))$ of $\pi_1(K) \simeq \pi$. By definition, the fundamental group of \overline{K} is isomorphic to $\psi(\pi_1(X))$. The map $\Psi : X \rightarrow K$ lifts to a map $\overline{\Psi} : X \rightarrow \overline{K}$ uniquely defined up to deck transformations such that $\Psi = p \circ \overline{\Psi}$. The index $[\pi : \psi(\pi_1(X))]$ of $\psi(\pi_1(X))$ in π is finite, otherwise \overline{K} is noncompact. In this case, the homology group $H_n(\overline{K}; \mathbb{Z})$ vanishes and $\Psi_*[X] = p_*(\overline{\Psi}_*[X])$ is trivial in homology, hence a contradiction. Thus, $|\deg p| = [\pi : \psi(\pi_1(X))]$. Since $\deg \Psi = \deg p \cdot \deg \overline{\Psi} = 1$, the index of $\psi(\pi_1(X))$ in π equals 1. Therefore, $\psi(\pi_1(X)) = \pi$. Hence ψ is surjective and the same goes for ϕ . \square

The following example shows that the surjectivity of ψ does not hold anymore when K is not an n -dimensional manifold (of course, here, we replace $[K]$ by $\Phi_*[M]$).

Example 3.8. Consider the connected sum $M = (S^1 \times S^{n-1}) \# \mathbb{T}^n$ with $n \geq 3$ (we can replace the n -torus \mathbb{T}^n by any essential manifold) and a $K(\pi_1(M), 1)$ -space K . Here, $\pi_1(M) \simeq \mathbb{Z} * \pi_1(\mathbb{T}^n) \simeq \mathbb{Z} * \mathbb{Z}^n$ from Van Kampen's theorem. Since K is aspherical, the canonical map $\Phi : M \rightarrow K$ induced by the identity homomorphism on $\pi_1(M)$ factorizes through $\Phi : M \rightarrow (S^1 \times S^{n-1}) \vee \mathbb{T}^n \xrightarrow{i \vee j} K$. Furthermore, the map $j : \mathbb{T}^n \rightarrow K$ represents $\Phi_*[M]$ and $j_* : \pi_1(\mathbb{T}^n) \rightarrow \mathbb{Z} * \pi_1(\mathbb{T}^n)$ is not surjective.

One can kill the kernel of ψ by attaching 2-cells at X . More precisely, we have the following.

Lemma 3.9. *There exists an H -extension X' of X and a map $\Psi' : X' \rightarrow K$ of degree 1 such that the homomorphism $\Psi'_* : \pi_1(X') \rightarrow \pi$ induced by Ψ' is an isomorphism which agrees with ψ' defined after (3.5).*

Further, since X' is an H -extension of X , $\text{MinEnt}(X') = \text{MinEnt}_H(X)$.

Proof. Note that the groups $\pi_1(X)$ and π have finite presentations. The kernel H of the epimorphism $\psi : \pi_1(X) \rightarrow \pi$ is generated by a finite number of elements. Define an H -extension X' of X by attaching 2-cells along a finite number of (simplicial) loops representing a finite generating set of H . By construction, the map Ψ extends into a map $\Psi' : X' \rightarrow K$ of degree 1 such that the homomorphism induced between the fundamental groups is an isomorphism which agrees with ψ' . \square

Similar arguments lead to the following (see also [4, Lemme 3.11]).

Lemma 3.10. *There exists an oriented closed n -dimensional manifold M' and a map $\Phi' : M' \rightarrow K$ of degree 1 such that the homomorphism $\phi' : \pi_1(M') \rightarrow \pi$ induced by Φ' is an isomorphism and $\text{MinEnt}(M') = \text{MinEnt}_G(M)$.*

Proof. The kernel G of the epimorphism $\phi : \pi_1(M) \rightarrow \pi$ is generated by a finite number of elements represented by disjoint simple loops $\gamma_1, \dots, \gamma_k$ of M . Consider disjoint tubular neighborhoods of the γ_i 's homeomorphic to $S^1 \times B^{n-1}$. Remove these neighborhoods from M and glue $D_i^2 \times S^{n-2}$ along their boundaries $S_i^1 \times S^{n-2}$ where D_i^2 is a disk bounding S_i^1 . The oriented closed n -manifold so-obtained is noted M' .

Since K is aspherical and the images by Φ of the loops γ_i are contractible, the map Φ induces a map $\Phi' : M' \rightarrow K$ which represents the fundamental class of K . By construction, the homomorphism $\phi' : \pi_1(M') \rightarrow \pi$ induced by Φ' is an isomorphism.

Let Y be the extension of M' obtained by attaching an $(n-1)$ -cell B_i^{n-1} along a copy S^{n-2} of $D_i^2 \times S^{n-2}$ for every $1 \leq i \leq k$. This

ensures us that the spheres S^{n-2} of $D_i^2 \times S^{n-2}$ are contractible in Y . Therefore, there exists an n -monotone map $f : M \longrightarrow Y$ of degree 1 such that $\phi = \phi' \circ f_*$. From Lemma 3.5 and Lemma 3.6, we have $\text{MinEnt}_G(M) \leq \text{MinEnt}(Y) = \text{MinEnt}(M')$.

The reverse inequality, which will not be used in the sequel, can be obtained as follows. Let Z be the G -extension of M obtained by attaching 2-cells D_i^2 along the γ_i 's. The projections of the second factor of $D_i^2 \times S^{n-2}$ onto D_i^2 give rise to an n -monotone map of degree 1 from M' onto Z which induces an isomorphism between the fundamental groups. As previously, from Lemma 3.5 and Lemma 3.6, we obtain $\text{MinEnt}(M') \leq \text{MinEnt}(Z) = \text{MinEnt}_G(M)$. \square

The following result rests on a construction of [4, Théorème 3.7]. The restriction on the dimension, i.e., $n \neq 3$, is required at this point precisely (see Remark 3.12).

Lemma 3.11. *There exists an extension X'' of X' and a degree 1 map $\Phi'' : M' \longrightarrow X''$ such that the homomorphism $\phi'' : \pi_1(M') \longrightarrow \pi_1(X'')$ induced by Φ'' is an isomorphism which agrees with $\psi'^{-1} \circ \phi'$ and*

$$\text{MinEnt}(X'') \geq \text{MinEnt}(M'). \quad (3.8)$$

Further, since X'' is an extension of X' , $\text{MinEnt}(X'') = \text{MinEnt}(X')$.

Proof. By assumption, $\psi' : \pi_1(X') \longrightarrow \pi$ and $\phi' : \pi_1(M') \longrightarrow \pi$ are isomorphisms and $\Psi'_*[X'] = \Phi'_*[M'] = [K]$ in $H_n(K; \mathbb{Z}) \simeq H_n(\pi; \mathbb{Z})$. We can apply the theorem 3.7 of [4] which yields an extension X'' of X' and a map $\Phi'' : M' \longrightarrow X''$ such that the homomorphism $\phi'' : \pi_1(M') \longrightarrow \pi_1(X'')$ induced by Φ'' satisfies $\phi'' = \psi'^{-1} \circ \phi'$.

Remark 3.12. Strictly speaking, this result has been stated for two manifolds M_1 and M_2 (see [4, Théorème 3.7] for the notations). Furthermore, it does not hold anymore when M_2 is replaced by a polyhedron (see [4] for a counterexample). However, when M_1 is a pseudomanifold and M_2 is a manifold, the proof still applies which permits us to conclude.

Let us develop this point by recalling the steps of the construction of [4, Théorème 3.7]:

- 1) The homotopy groups of M_1 of dimension i with $2 \leq i \leq n-1$ can be successively killed by attaching cells of dimensions $i+1$. The resulting spaces are CW-complexes noted $M_1(i+1)$ with $M_1(n)$ homotopically equivalent to the n -skeleton $K(\pi, 1)^{(n)}$ of $K(\pi, 1)$.
- 2) The cellular approximation theorem applied to the classifying map of M_2 yields a map $g : M_2 \longrightarrow M_1(n)$. Note that $M_1(n)$ is not an extension of M_1 .

- 3) The map g is not uniquely defined up to homotopy. However, it may be chosen so that it can be deformed into a map $\hat{g} : M_2 \longrightarrow M_1(n-1)$ inducing the same homomorphism as g between the fundamental groups.
- 4) By compactness, the map \hat{g} can be deformed into a map whose image is a finite subcomplex of $M_1(n-1)$ and so an extension of M_1 .

Step 3) is the main step of the construction. It relies on the obstruction to extend a map inducing a fundamental group isomorphism. When such an obstruction is trivial, the extension can be performed without modifying the original map on its 2-codimensional skeleton and, in particular, on its 2-skeleton when $n \geq 4$. This ensures that the extension still induces an isomorphism between the fundamental groups.

The proof of 3) requires that M_2 is a manifold to apply the Poincaré duality. However, the assumption that M_1 is a manifold plays no role in any of the four steps of the construction (M_1 is turned into a CW-complex from the first step by attaching cells to it). For the purpose of the proof, it is enough that M_1 is a pseudomanifold.

Let us now conclude the proof of Lemma 3.11. Since K is aspherical, the map Ψ' extends to X'' into a degree 1 map. This map, still denoted Ψ' , induces a homomorphism between the fundamental groups which agrees with $\psi' : \pi_1(X'') \simeq \pi_1(X') \longrightarrow \pi$. The homomorphism $\psi' \circ \phi''$ agrees with ϕ' . Therefore, since K is aspherical, the map $\Psi' \circ \Phi''$ is homotopic to the degree 1 map Φ' . This shows that $\deg \Psi' \circ \Phi'' = \deg \Psi' \cdot \deg \Phi'' = 1$. Thus, the degree of Φ'' is equal to 1.

The polyhedron X'' is an H -extension of the closed n -pseudomanifold X . Therefore, it admits a cellular decomposition with exactly one n -cell (see Lemma 2.2). As noticed in [4, Propriété 3.12], the proof of [15, Theorem 4.1] still applies in this case. This implies from [4, Propriété 3.12] that the map $\Phi'' : M' \longrightarrow X''$ of degree 1 is homotopic to an n -monotone map $\bar{\Phi}'' : M' \longrightarrow X''$. We deduce from Lemma 3.5 that $\text{MinEnt}(X'') \geq \text{MinEnt}(M')$. \square

The three previous lemmas show that every geometric cycle X representing $\Phi_*[M]$ satisfies

$$\text{MinEnt}_G(M) \leq \text{MinEnt}_H(X). \quad (3.9)$$

Thus, $\text{MinEnt}_G(M) \leq \text{MinEnt}(\Phi_*[M])$.

The reverse inequality is obvious. Therefore, Theorem 3.1 is proved for $n \geq 4$.

When $n = 2$, M is a surface of genus $\gamma \geq 1$. In this case, K is homeomorphic to a surface of genus at most γ . Let $\Psi : X \rightarrow K$ be a geometric cycle representing $\Phi_*[M] = [K]$ in homology. From the definition of pseudomanifolds, there exists a finite number of points x_1, \dots, x_k of X such that $X \setminus \{x_1, \dots, x_k\}$ is a two-dimensional manifold and every point x_i admits a neighborhood homeomorphic to a cone over a finite number of circles. The completed space of $X \setminus \{x_1, \dots, x_k\}$ is a closed orientable surface N . The inclusion $X \setminus \{x_1, \dots, x_k\} \hookrightarrow X$ extends into a 2-monotone map $j : N \rightarrow X$ such that $j_*[N] = [X]$. Thus, the map $\Psi_0 = \Psi \circ j : N \rightarrow K$ represents $\Phi_*[M] = [K]$. We can assume that the genus of K is greater than one, otherwise the result is obvious since $\pi_1(M)/G$ is isomorphic to \mathbb{Z}^2 and the minimal G -entropy of M is then zero. The degree 1 map Ψ_0 induces a homomorphism $\psi_0 : \pi_1(N) \rightarrow \pi$ between the fundamental groups, whose kernel is noted H_0 . Arguing as in [11, Théorème 8.1], one can show that $\text{MinEnt}_{H_0}(N) = \text{MinEnt}(M)$. Since j is 2-monotone and $\psi_0 = \psi \circ j_*$, we deduce that $\text{MinEnt}_{H_0}(N) \leq \text{MinEnt}_H(X)$ from Lemma 3.5. Therefore, $\text{MinEnt}(M) \leq \text{MinEnt}_H(X)$. As previously, Theorem 3.1 immediately follows when $n = 2$.

4. ENTROPY OF EXTREMAL GEOMETRIC CYCLES

We introduce in this section the notion of regular geometric cycles and bound from above their relative entropy in terms of their relative systolic volume.

In dimension greater than two, we do not know whether the infimum in inequality (1.1) is achieved by any metric. The likely absence of extremal metrics in general makes the study of systolic geometry delicate. In order to get around this difficulty M. Gromov proved that every essential manifold can be represented by a regular geometric cycle (see below). These regular geometric cycles share enough patterns we expect from possible extremal metrics to play the role of substitute in their absence. Note however that regular geometric cycles are not necessarily homeomorphic to the manifold they represent, neither do they necessarily have the same fundamental group.

Given a discrete group π , the optimal systolic volume of a homology class $h \in H_n(\pi; \mathbb{Z})$ is defined as

$$\sigma(h) = \inf_X \sigma_\psi(X) \quad (4.1)$$

where $X = (X, \Psi, g)$ runs over all the geometric cycles representing h .

M. Gromov showed in [18, §6] that there exists a positive constant c_n such that every nontrivial homology class $h \in H_n(\pi; \mathbb{Z})$ satisfies

$$\sigma(h) \geq c_n \quad (4.2)$$

Furthermore, he proved in [18, p. 71] that there exists $c_n > 0$ such that, for every $\varepsilon > 0$, every nontrivial homology class $h \in H_n(\pi; \mathbb{Z})$ can be represented by a geometric cycle $X = (X, \Psi, g)$ such that

$$\sigma_\psi(X, g) \leq (1 + \varepsilon)\sigma(h) \quad (4.3)$$

and

$$\text{vol}_g(B(x, R)) \geq c_n R^n \quad (4.4)$$

for every $x \in X$ and $\varepsilon \leq R/\text{sys}_\psi(X, g) \leq \frac{1}{2}$, where $\text{vol}_g(B(x, R))$ is the volume of the g -ball centered at x with radius R in X . In (4.4), we can take $c_n = n^{-n+2}((n-1)!\sqrt{n!})^{-n+1}$.

Such geometric cycles are said to be ε -regular.

Arguing as in [26] (see also [22]), we show the following.

Proposition 4.1. *Every ε -regular geometric cycle (X, Ψ, g) representing a nontrivial homology class $h \in H_n(\pi; \mathbb{Z})$ satisfies*

$$\text{Ent}_H(X, g) \leq \frac{1}{\beta \text{sys}_\psi(X, g)} \log \left(\frac{\sigma_\psi(X, g)}{c_n \alpha^n} \right), \quad (4.5)$$

whenever $\alpha \geq \varepsilon$, $\beta > 0$ and $4\alpha + \beta < \frac{1}{2}$, where c_n is given by (4.4).

Proof. Let $x_0 \in X$ be a fixed basepoint. Consider a maximal system of disjoint balls

$$B_i = B(x_i, R) \subset X \quad (4.6)$$

of radius $R = \alpha \text{sys}_\psi(X, g)$ and centers x_i with $i \in I$, including x_0 . Since the geometric cycle (X, Ψ, g) is ε -regular, we have

$$\text{vol } B_i \geq c_n \alpha^n \text{sys}_\psi(X, g)^n \quad \forall i \in I. \quad (4.7)$$

Therefore, this system admits at most $\frac{\text{vol}(X, g)}{c_n \alpha^n \text{sys}_\psi(X, g)^n}$ balls. Thus,

$$|I| \leq \frac{\sigma_\psi(X, g)}{c_n \alpha^n}. \quad (4.8)$$

Let $c : [0, T] \rightarrow X$ be a geodesic loop of length T based at x_0 whose image by Ψ is noncontractible. Let

$$m = \left\lfloor \frac{T}{\beta \text{sys}_\psi(X, g)} \right\rfloor \quad (4.9)$$

be the integer part. The point $p_0 = x_0$, together with the points

$$p_k = c(k\beta \text{sys}_\psi(X, g)), \quad k = 1, \dots, m$$

and the point $p_{m+1} = x_0$, partition the loop c into $m + 1$ segments of length at most $\beta \text{sys}_\psi(X, g)$. Since the system of balls B_i is maximal, the disks of radius $2R = 2\alpha \text{sys}_\psi(X, g)$ centered at x_i cover X . Therefore, for every p_k , a nearest point q_k among the centers x_i , is at distance at most $2R$ from p_k . Consider the loop

$$\alpha_k = c_k \cup [p_{k+1}, q_{k+1}] \cup [q_{k+1}, q_k] \cup [q_k, p_k],$$

where c_k is the arc of c joining p_k to p_{k+1} , while $[x, y]$ denotes a minimizing path joining x to y . Then

$$\text{length}(\alpha_k) \leq 2(4\alpha + \beta) \text{sys}_\psi(X, g) < \text{sys}_\psi(X, g),$$

by our hypothesis on α, β . Thus the image by Ψ of the loops α_k is contractible. The same is true for the loops $c_0 \cup [p_1, q_1] \cup [q_1, x_0]$ and $c_m \cup [x_m, q_m] \cup [q_m, p_m]$.

Therefore, the image by Ψ of the geodesic loop c is homotopic to the image by Ψ of a piecewise geodesic loop $c' = (x_0, q_1, \dots, q_m, x_0)$. Note that since the length of minimizing paths from p_k to q_k is at most $(4\alpha + \beta) \text{sys}_\psi(X, g) < \frac{1}{2} \text{sys}_\psi(X, g)$, their images by Ψ are homotopic with fixed endpoints. Thus, the choice of a minimizing path between p_k and q_k does not matter.

Thus, c and c' represent the same class in $\text{Aut}(X_H) \simeq \pi_1(X)/H$ where $H = \ker \psi$. Furthermore, two closed geodesics c_1 and c_2 which induce two distinct classes in $\text{Aut}(X_H)$ give rise to two loops c'_1 and c'_2 with the same property. Thus, the number $P_H(T)$ of classes in $\text{Aut}(X_H)$ which can be represented by loops of length at most T based at x_0 satisfies

$$\begin{aligned} P_H(T) &\leq |I|^m \\ &\leq |I|^{\frac{T}{\beta \text{sys}_\psi(X, g)}} \\ &\leq \left(\frac{\sigma_\psi(X, g)}{c_n \alpha^n} \right)^{\frac{T}{\beta \text{sys}_\psi(X, g)}}, \end{aligned} \tag{4.10}$$

and the proposition now follows from Lemma 2.3. \square

5. RELATIVE SYSTOLIC VOLUME AND RELATIVE MINIMAL ENTROPY

Using the results established in the previous sections, we show that under some topological conditions, the relative systolic volume of manifolds is bounded from below in terms of their relative minimal entropy. More precisely, we have the following.

Theorem 5.1. *Let π be a discrete group. Let $\Phi : M \longrightarrow K$ be a degree 1 map between two oriented closed manifolds of the same dimension $n \neq 3$*

such that K is a $K(\pi, 1)$ -space. Then, there exists a positive constant C_n depending only on n such that

$$\sigma_\phi(M) \geq C_n \frac{\text{MinEnt}_G(M)^n}{\log^n(1 + \text{MinEnt}_G(M))}. \quad (5.1)$$

where G is the kernel of the homomorphism $\phi : \pi_1(M) \longrightarrow \pi$ induced by Φ .

Remark 5.2. Since $\sigma(M) \geq \sigma_\phi(M)$, we can replace $\sigma_\phi(M)$ by $\sigma(M)$ in inequality (5.1).

Example 5.3. Recall M. Gromov's isosystolic inequality for surfaces of large genus (see [18, 6.4.D'], [26] and [6]): there exists a positive constant c such that every closed surface Σ_γ of genus $\gamma > 1$ satisfies

$$\sigma(\Sigma_\gamma) \geq c \frac{\gamma}{(\log \gamma)^2}.$$

Theorem 5.1 generalizes this result for surfaces of large genus with few small handles. More precisely, consider a degree 1 map $\Phi : \Sigma_\gamma \longrightarrow \Sigma_{\gamma'}$ between two closed surfaces of genus γ and $\gamma' > 1$. From Remark 3.2 and inequality (1.9), we have

$$\text{MinEnt}_G(\Sigma_\gamma) = \text{MinEnt}(\Sigma_{\gamma'}) = 4\pi(\gamma' - 1).$$

This implies that

$$\sigma_\phi(\Sigma_\gamma) \geq c' \frac{\gamma'}{(\log \gamma')^2}$$

for some $c' > 0$. Thus, the area of a closed Riemannian surface of large genus with, let's say, a single "small" handle is still large.

Proof of Theorem 5.1. We will use the same notations as in the previous sections. From Proposition 4.1, every ε -regular cycle (X, Ψ, g) representing the image $\Phi_*[M]$ of the fundamental class of M in $H_n(K; \mathbb{Z}) \simeq H_n(\pi; \mathbb{Z})$ satisfies

$$\text{Ent}_H(X, g)^n \text{vol}(X, g) \leq \frac{\sigma_\psi(X, g)}{\beta^n} \log^n \left(\frac{\sigma_\psi(X, g)}{c_n \alpha^n} \right) \quad (5.2)$$

whenever $\alpha \geq \varepsilon$, $\beta > 0$ and $4\alpha + \beta < \frac{1}{2}$, with $\sigma_\psi(X, g) \leq (1 + \varepsilon)\sigma_\phi(M)$. From Theorem 3.1, we have

$$\text{MinEnt}_G(M)^n = \text{MinEnt}(\Phi_*[M])^n \leq \text{Ent}_H(X, g)^n \text{vol}(X, g). \quad (5.3)$$

Therefore, we deduce that

$$\text{MinEnt}_G(M)^n \leq \frac{\sigma_\phi(M)}{\beta^n} \log^n \left(\frac{\sigma_\phi(M)}{c_n \alpha^n} \right) \quad (5.4)$$

whenever $\alpha, \beta > 0$ and $4\alpha + \beta \leq \frac{1}{2}$.

Define

$$\alpha = \begin{cases} \frac{1}{10} & \text{if } n^n c_n \leq 1 \\ \frac{1}{10n \sqrt[n]{c_n}} & \text{otherwise} \end{cases} \quad \text{and} \quad \beta = \begin{cases} \frac{1}{10} n \sqrt[n]{c_n} & \text{if } n^n c_n \leq 1 \\ \frac{1}{10} & \text{otherwise.} \end{cases}$$

In both cases, we have $4\alpha + \beta \leq \frac{1}{2}$ and $\beta = n\alpha \sqrt[n]{c_n}$.

Let $\rho = \left(\frac{\sigma_\phi(M)}{c_n \alpha^n} \right)^{\frac{1}{n}}$ and $\delta = \frac{\beta}{n\alpha \sqrt[n]{c_n}} \text{MinEnt}_G(M)$. For our choices of α and β , we have $\delta = \text{MinEnt}_G(M)$. Note also that $\rho \geq 1$ since $\sigma_\phi(M) \geq \frac{c_n}{2^n}$ from inequality (4.4).

For $\delta \geq e$, inequality (5.4) yields the following estimate:

$$\begin{aligned} \rho \log \rho &\geq \delta \\ &\geq \delta - \frac{\delta \log \log \delta}{\log \delta} \\ &= \frac{\delta}{\log \delta} \log \left(\frac{\delta}{\log \delta} \right). \end{aligned}$$

The function $x \log x$ is increasing for $x \geq \frac{1}{e}$. Thus, since $\rho \geq 1$ and $\frac{\delta}{\log \delta} \geq e$ for $\delta \geq e$, we deduce that $\rho \geq \frac{\delta}{\log \delta} \geq \frac{\delta}{\log(1+\delta)}$ for $\delta \geq e$.

For $\delta \leq e$, we have $0 \leq \frac{\delta}{\log(1+\delta)} \leq \frac{e}{\log(1+e)}$. Therefore, we obtain $\rho \geq 1 \geq \frac{\log(1+e)}{e} \frac{\delta}{\log(1+\delta)}$ for $\delta \leq e$.

In conclusion, for every $\delta \geq 0$, the following holds:

$$\rho \geq \frac{\log(1+e)}{e} \frac{\delta}{\log(1+\delta)}. \quad (5.5)$$

Hence,

$$\sigma_\phi(M) \geq c_n \alpha^n \frac{\log^n(1+e)}{e^n} \frac{\text{MinEnt}_G(M)^n}{\log^n(1 + \text{MinEnt}_G(M))}. \quad (5.6)$$

Therefore, inequality (5.1) holds with $C_n = c_n \alpha^n \frac{\log^n(1+e)}{e^n}$. \square

6. RELATIVE SYSTOLIC VOLUME AND ALGEBRAIC ENTROPY

In this section, we show how the relative entropy of a polyhedron is related to the algebraic entropy of a quotient of its fundamental group. Then, we derive a lower bound on the relative systolic volume of some manifolds in terms of the algebraic entropy of some groups. Specifically, we prove the following.

Theorem 6.1. *Let π be a discrete group. Let $\Phi : M \longrightarrow K$ be a degree 1 map between two oriented closed manifolds of the same dimension n such*

that K is a $K(\pi, 1)$ -space. Then, there exists a positive constant C_n depending only on n such that

$$\sigma_\phi(M) \geq C_n \frac{\text{Ent}_{alg}(\pi)}{\log(1 + \text{Ent}_{alg}(\pi))} \quad (6.1)$$

where $\phi : \pi_1(M) \longrightarrow \pi$ is the homomorphism induced by Φ .

Remark 6.2. We can replace $\sigma_\phi(M)$ by $\sigma(M)$ in inequality (6.1). Furthermore, the integer n can be equal to 3.

Remark 6.3. Let F_k be the free group with k generators, $\pi_1(\Sigma_\gamma)$ be the fundamental group of a closed surface of genus γ and π be a discrete group with k generators and p relations. Then, we have

- $\text{Ent}_{alg}(F_k) \geq \log(2k - 1)$ (see [20, 5.13] or [21, VII.B]);
- $\text{Ent}_{alg}(\pi_1(\Sigma_\gamma)) \geq \log(4\gamma - 3)$ (see [21, VII.B]);
- $\text{Ent}_{alg}(\pi) \geq c \log(2(k - p) - 1)$ for some $c > 0$ (see [20, 5.14] and [17, p. 82-83]).

As shown in the following example, Theorem 6.1 may yield nontrivial lower bounds on the systolic volume when the contents of Theorem 1.2 and Theorem 1.3 are trivial.

Example 6.4. Consider the closed aspherical manifolds

$$M_\gamma = \Sigma_\gamma \times \mathbb{T}^{n-2}$$

of dimension $n \geq 3$, where Σ_γ is the closed surface of genus γ and \mathbb{T}^{n-2} is the $(n - 2)$ -torus. The group $\pi_1(\Sigma_\gamma)$ is a quotient of $\pi_1(M)$. Therefore,

$$\text{Ent}_{alg}(\pi_1(M_\gamma)) \geq \text{Ent}_{alg}(\Sigma_\gamma) \geq \log(4\gamma - 3).$$

From Theorem 6.1, we obtain

$$\sigma(M_\gamma) \gtrsim \frac{\log \gamma}{\log \log \gamma},$$

which goes to infinity as $\gamma \rightarrow \infty$. Such an estimate follows neither from Theorem 1.2 nor from Theorem 1.3 since $\text{MinEnt}(M_\gamma) = ||M_\gamma|| = 0$.

In this second example, we show that the conclusion of Theorem 6.1 does not hold without some topological conditions on M .

Example 6.5. Fix a closed aspherical manifold M of dimension $n \geq 3$. Consider the sequence of essential manifolds

$$M_k = M \# S^1 \times S^{n-1} \# \dots \# S^1 \times S^{n-1}$$

where we take the connected sum of M with k copies of $S^1 \times S^{n-1}$. As shown in [5], adding handles to a manifold does change the value of the optimal systolic volume. Thus, for every k , we have

$$\sigma(M_k) = \sigma(M).$$

On the other hand, the difference between the number of generators and the number of relations of $\pi_1(M_k)$ is equivalent to k . Therefore,

$$\text{Ent}_{alg}(\pi_1(M_k)) \gtrsim \log k.$$

This shows that inequality (6.1) does not hold in general: some topological condition must be satisfied as in Theorem 6.1.

The following result is a relative version of [20, Theorem 5.16] (which is incorrectly stated).

Lemma 6.6. *Let X be a closed polyhedron and H be a normal subgroup of the fundamental group $\pi_1(X)$ of X . Then, every piecewise linear Riemannian metric g on X satisfies*

$$\text{Ent}_{alg}(\pi_1(X)/H) \leq 2 \text{Diam}(X, g) \text{Ent}_H(X, g). \quad (6.2)$$

Proof. Fix x_0 in X . The fundamental group $\pi_1(X, x_0)$ of X is generated by a finite set Σ formed of classes represented by loops based at x_0 of length at most $2 \text{Diam}(X, g)$ (see [20, 3.22]). Denote by $\bar{\Sigma}$ the generating set of $\Gamma := \pi_1(X, x_0)/H$ induced by Σ . Every class of $|\cdot|_{\bar{\Sigma}}$ -length at most k can be represented by a loop based at x_0 of length at most $2k \text{Diam}(X, g)$. Therefore, $N_{\bar{\Sigma}}(k) \leq P_H(2k \text{Diam}(X, g))$ where $N_{\bar{\Sigma}}$ and P_H are defined in (1.11) and Lemma 2.3. Thus,

$$\frac{\log(N_{\bar{\Sigma}}(k))}{k} \leq 2 \text{Diam}(X, g) \frac{\log(P_H(2k \text{Diam}(X, g)))}{2k \text{Diam}(X, g)}. \quad (6.3)$$

From Lemma 2.3, we obtain

$$\text{Ent}_{alg}(\Gamma) \leq \text{Ent}_{alg}(\Gamma, \bar{\Sigma}) \leq 2 \text{Diam}(X, g) \text{Ent}_H(X, g). \quad (6.4)$$

when k goes to infinity. \square

We can now prove the main result of this section.

Proof of Theorem 6.1. Every ε -regular geometric cycle (X, Ψ, g) representing the image $\Phi_*[M]$ of the fundamental class of M admits at least $\left\lceil \frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \right\rceil$ disjoint balls of radius $\frac{1}{2} \text{sys}_\psi(X, g)$. Combined with the inequality (4.4), this yields

$$\text{vol}(X, g) \geq \left\lceil \frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \right\rceil \frac{c_n}{2^n} \text{sys}_\psi(X, g)^n. \quad (6.5)$$

Hence,

$$\left\lceil \frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \right\rceil \leq \frac{2^n}{c_n} \sigma_\psi(X, g) \quad (6.6)$$

From Lemma 3.7 (which holds for every integer n), we have $\pi_1(X)/H \simeq \pi$. Now, Lemma 6.6 and Proposition 4.1 imply that

$$\text{Ent}_{alg}(\pi) \leq 2 \text{Diam}(X, g) \text{Ent}_H(X, g) \quad (6.7)$$

$$\leq \frac{2 \text{Diam}(X, g)}{\beta \text{sys}_\psi(X, g)} \log \left(\frac{\sigma_\psi(X, g)}{c_n \alpha^n} \right) \quad (6.8)$$

whenever $\alpha, \beta > 0$ and $4\alpha + \beta < \frac{1}{2}$.

If $\frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \geq 2$, then $\frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \leq 2 \left\lceil \frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \right\rceil$.

Otherwise, $\frac{\text{Diam}(X, g)}{\text{sys}_\psi(X, g)} \leq \frac{2^{n+1}}{c_n} \sigma_\psi(X, g)$ since $\sigma_\psi(X, g) \geq \frac{c_n}{2^n}$ from (4.4).

In both case, we deduce from (6.6) and (6.8) that

$$\text{Ent}_{alg}(\pi) \leq \frac{2^{n+1}}{\beta c_n} \sigma_\psi(X, g) \log \left(\frac{\sigma_\psi(X, g)}{c_n \alpha^n} \right) \quad (6.9)$$

whenever $\alpha, \beta > 0$ and $4\alpha + \beta < \frac{1}{2}$, with $\sigma_\psi(X, g) \leq (1 + \varepsilon) \sigma_\phi(M)$.

Therefore, we have

$$\text{Ent}_{alg}(\pi) \leq \frac{2^{n+1}}{\beta c_n} \sigma_\phi(M) \log \left(\frac{\sigma_\phi(M)}{c_n \alpha^n} \right) \quad (6.10)$$

whenever $\alpha, \beta > 0$ and $4\alpha + \beta \leq \frac{1}{2}$.

Arguing as in the proof of Theorem 5.1, we obtain

$$\sigma_\phi(M) \geq c_n \alpha^n \frac{\log(1 + e)}{e} \frac{\text{Ent}_{alg}(\pi)}{\log(1 + \text{Ent}_{alg}(\pi))} \quad (6.11)$$

with $\alpha = \frac{1}{10}$. Therefore, inequality (6.1) holds with $C_n = c_n \alpha^n \frac{\log(1+e)}{e}$. \square

REFERENCES

- [1] Babenko, I.: Asymptotic invariants of smooth manifolds, *Russian Acad. Sci. Izv. Math.* **41** (1993) 1–38, MR1208148, Zbl 0812.57022.
- [2] Babenko, I.: Forte souplesse intersystolique de variétés fermées et de polyèdres. *Annales de l'Institut Fourier* **52** (2002) 1259–1284, MR2169832, Zbl 1023.53025.
- [3] Babenko, I.: Géométrie systolique des variétés de groupe fondamental \mathbb{Z}_2 . *Sémin. Théor. Spectr. Géom. Grenoble* **22** (2004) 25–52, MR2136133, Zbl pre02149962.
- [4] Babenko, I.: Nature topologique des systoles unidimensionnelles. Preprint available at <http://www.math.univ-montp2.fr/prepublications/03-07.pdf>
- [5] Babenko, I.; Balacheff, F.: Géométrie systolique des sommes connexes et des revêtements cycliques. *Math. Ann.* **333** (2005) 157–180, MR2169832, Zbl pre02235442.
- [6] Balacheff, F.: Sur des problèmes de la géométrie systolique. *Sémin. Théor. Spectr. Géom. Grenoble* **22** (2004) 71–82, MR2136136, Zbl pre02149965.

- [7] Bangert, V.; Croke, C.; Ivanov, S.; Katz, M.: Boundary case of equality in optimal Loewner-type inequalities, *Trans. A.M.S.*, to appear.
- [8] Bangert, V.; Katz, M.: Stable systolic inequalities and cohomology products, *Comm. Pure Appl. Math.* **56** (2003), 979-997, MR1990484, Zbl 1038.53031.
- [9] Bangert, V.; Katz, M.: An optimal Loewner-type systolic inequality and harmonic one-forms of constant norm. *Comm. Anal. Geom.* **12** (2004), 701-730, MR2128608, Zbl 1068.53027.
- [10] Bavard, C.: Inégalité isosystolique pour la bouteille de Klein, *Math. Ann.* **274** (1986), 439-441, MR0960206, Zbl 0578.53032.
- [11] Besson, G.; Courtois, G.; Gallot, S.: Volume et entropie minimale des espaces localement symétriques. *Invent. Math.* **103** (1991), 417-445, MR1085114, Zbl 0723.53029.
- [12] Berger, M.: Systoles et applications selon Gromov, Sémin. Bourbaki, Astérisque **216** (1993) 279-310, MR1246401, Zbl 0789.53040.
- [13] Buser, P.; Sarnak, P.: On the period matrix of a Riemann surface of large genus. With an appendix by J. H. Conway and N. J. A. Sloane. *Invent. Math.* **117** (1994), 27-56, MR1269424, Zbl 0814.14033.
- [14] Croke, C.; Katz, M.: Universal volume bounds in Riemannian manifolds, Yau, S.-T. (ed.), *Surveys in Differential Geometry VIII*, Lectures on Geometry and Topology held in honor of Calabi, Lawson, Siu, and Uhlenbeck at Harvard University, May 3-5, 2002. Somerville, MA: International Press, 2003, pp. 109-137, MR2039987, Zbl 1051.53026.
- [15] Epstein, D. B. A.: The degree of a map, *Proc. London Math. Soc.* **16** (1966), 369-383, MR0192475, Zbl 0148.43103.
- [16] Freedman, M.: \mathbb{Z}_2 -systolic-freedom, Proceedings of the Kirbyfest (Berkeley, CA, 1998), 113-123, Geom. Topol. Monogr., vol. 2, Geom. Topol., Coventry, 1999. Available at <http://www.maths.warwick.ac.uk/gt/GTMon2/paper6.abs.html>, MR1734404, Zbl 1024.53029.
- [17] Gromov, M.: Volume and bounded cohomology, *Publ. IHES.* **56** (1981), 213-307, MR0686042, Zbl 0516.53046.
- [18] Gromov, M.: Filling Riemannian manifolds, *J. Differential Geom.* **18** (1983), 1-147, MR0697984, Zbl 0515.53037.
- [19] Gromov, M.: Systoles and intersystolic inequalities. In: *Actes de la Table Ronde de Géométrie Différentielle (Luminy 1992)*, 291-362. Séminaires et Congrès, **1**. Soc. Math. France, Institut Henri Poincaré, Paris 1996, MR1427763, Zbl 0877.53002.
- [20] Gromov, M.: Metric structures for Riemannian and non-Riemannian spaces. Progress in Mathematics **152**. Birkhäuser, Boston 1999, MR1699320, Zbl 0953.53002.
- [21] de la Harpe, P.: Topics in geometric group theory, Chicago Lectures in Mathematics, Univ. Chicago Press, 2000, MR1786869, Zbl 0965.20025.
- [22] Katok, A.: Entropy and closed geodesics, *Ergo. Th. Dynam. Sys.* **2** (1983), 339-365, MR0721728, Zbl 0525.58027.
- [23] Ivanov, S.; Katz, M.: Generalized degree and optimal Loewner-type inequalities, *Israel J. Math.* **141** (2004), 221-233, MR2063034, Zbl 1067.53031.
- [24] Katz, M.: Four-manifold systoles and surjectivity of period map, *Comment. Math. Helvetici* **78** (2003), 772-876, MR2016695, Zbl 1048.53031.

- [25] Katz, M.; Rudyak, Yu.: Lusternik-Schnirelmann category and systolic category of low dimensional manifolds, *Comm. Pure Appl. Math.*, to appear.
- [26] Katz, M.; Sabourau, S.: Entropy of systolically extremal surfaces and asymptotic bounds. *Ergo. Th. Dynam. Sys.* **25** (2005) 1209–1220, MR2158402, Zbl pre02234476.
- [27] Katz, M.; Suci A.: Systolic freedom of loop space, *Geom. Func. Anal.* **11** (2001), 60–73, MR1829642, Zbl 1048.53030.
- [28] Manning A.: Topological entropy for geodesic flows, *Ann. of Math.* (2) **110** (1979), 567–573, MR0554385, Zbl 0426.58016.
- [29] Pu, P. M.: Some inequalities in certain nonorientable Riemannian manifolds, *Pacific J. Math.* **2** (1952), 55–71, MR0048886, Zbl 0046.39902.
- [30] Sakai, T.: A proof of the isosystolic inequality for the Klein bottle, *Proc. Amer. Math. Soc.* **104** (1988), 589–590, MR0962833, Zbl 0692.53019.
- [31] Spanier, E.: Algebraic topology. Springer-Verlag, New-York, 1966, MR0210112, Zbl 0145.43303.

LABORATOIRE DE MATHÉMATIQUES ET PHYSIQUE THÉORIQUE, UNIVERSITÉ DE
TOURS, PARC DE GRANDMONT, 37400 TOURS, FRANCE

E-mail address: `sabourau@lmpt.univ-tours.fr`